

A Study to Assess the Effectiveness of Video Assisted Teaching Programme Regarding Preventive Method of Farmer's Lung Among the Community People at Ramnagar, Kanpur, UP

Prarthana Mahapatra¹, Adarsh Kumar², Akanksha Verma³, Abhyarthana Mahapatra⁴, Deepak Suwalka⁵

¹Lecturer, Community Health Nursing, Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

²Assistant Professor, Medical Surgical Nursing, Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

³Nursing Tutor, Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

⁴Nursing Tutor, Divine Institute of Medical Science and Research, Baripada, Odisha, India

⁵Professor, Mental Health Nursing, Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

Corresponding Author:

Prarthana Mahapatra, Lecturer, Community Health Nursing,
Faculty of Nursing, Rama University, Kanpur, Uttar Pradesh, India

E-mail:

prarthanamahapatra@gmail.com

GFPSS-International Journal of Multidisciplinary Research is a journal of open access. In this journal, we allow all types of articles to be distributed freely and accessible under the terms of the creative common attribution- non-commercial-share. This allows the authors, readers and scholars and general public to read, use and to develop non-commercially work, as long as appropriate credit is given and the newly developed work are licensed with similar terms.

How to cite this article: Mahapatra P, Kumar A, Verma A, Mahapatra A, Suwalka D. A Study to Assess the Effectiveness of Video Assisted Teaching Programme Regarding Preventive Method of Farmer's Lung Among the Community People at Ramnagar, Kanpur, UP. GFPSS-IJMR 2025; 6:09: 3115-3119

Submitted: 15 September 2025; **Accepted:** 30 September 2025; **Published:** 10 October 2025

Abstract

Introduction: Farmer's lung disease (FLD) is a form of hypersensitivity pneumonitis (HP) caused by inhaling microorganisms from hay or grain stored in conditions of high humidity in the agricultural workplace. It is probably underdiagnosed, especially in northern Spain, where climatic conditions favor the development of this disease. The clinical presentation may vary, differentiating the chronic (exposure to lower concentrations of the antigen over a longer period time) and the acute forms (after exposure to high concentrations of the antigen).

Materials & Methods: The present study adopted a **quantitative evaluative research approach** to assess the effectiveness of the intervention. A **quasi-experimental one group pre-test post-test research design** was employed to evaluate the impact of the educational intervention on the selected variables. The **population** for the study comprised **community people**, and the **sample** included community members residing in **Ramnagar, Kanpur, Uttar Pradesh**. The **sample size** was 200 determined based on feasibility, availability, and the inclusion criteria of the study participants.

Results: The result of the study was In pre-test out of 200 members (25%) were having inadequate knowledge, (70%) were having moderately adequate knowledge, (5%) were having adequate knowledge in pre-test. While in in post-test out of 200 of members of (8%) were having inadequate knowledge, (72%) were having moderately adequate knowledge, (20%) were having adequate knowledge in post-test.

Conclusion: The study aimed to evaluate the effectiveness of a **video-assisted teaching program** on knowledge regarding **preventive methods of Farmer's Lung** among community members in Kanpur, Uttar Pradesh. Data were collected and statistically analyzed based on the study objectives. Results revealed that pre-test knowledge was mostly **moderate (70%)**, while post-test findings showed a marked improvement with **20% having adequate knowledge**, indicating the effectiveness of the teaching program.

Keywords: Effectiveness; Video Assisted Teaching; Preventive Method

Introduction:

Farmer's lung is considered a type of hypersensitivity pneumonitis that was initially identified in the 1700s by Italian researcher Bernardino Ramazzini. He examined exposure risks across multiple professions and found that the disease process was most abundantly found in breeders and farmers who were exposed to organic dust. Interest and knowledge of the disease were rekindled in 1932 when a county tuberculosis officer identified 5 cases of acute respiratory failure seemingly caused by moldy hay during the springtime. Hypersensitivity pneumonitis, as we understand it today, encompasses a wide breadth of exposures that may illicit a delayed allergic response, specifically in the smaller airways. Farmers, in particular, are exposed to various agents in the fields of their workplace. These include organic agents, inorganic agents, gasses, methane, pesticides, fertilizers, and more. The syndrome has great variability in symptom severity and presentation due in large part to the variable exposure durations and innumerable offending agents. Spread between people is nonexistent, however, common exposure history can lead to similar disease states.

The spores that cause Farmer's Lung are not infectious. Instead, they trigger an allergic reaction. Allergic reactions are produced by the immune system, the body's defence system that normally protects against infectious diseases. The immune system reacts when special substances called "antigens" enter the body. Antigens are usually found in germs such as infectious bacteria or viruses. The immune system reacts to neutralize these germs and prevent infections. Antigens are also found in harmless materials such as mould spores. Sometimes the immune system will react against these by producing a harmful reaction (e.g., an allergic reaction) against an antigen attached to material like a spore that is otherwise harmless.

Farmer's Lung develops in two steps, as other allergic reactions do. The first time a person inhales a large amount of dust and spores from mouldy hay, the immune system often responds by producing specific antibodies against the antigens. Antibodies are chemicals that circulate in the blood and attack specific antigens. Once the body has produced these antibodies, there is a chance that further exposure to mouldy dust can generate a hypersensitive type of allergic reaction. This reaction is typical of Farmer's Lung.

Objectives

1. To assess the existing level of knowledge regarding preventive method of farmer's lung among the community people at Ramnagar, Kanpur, UP.
2. To evaluate the effectiveness of video assisted teaching programme regarding preventive method of farmer's lung among the community people at Ramnagar, Kanpur, UP. .
3. To find out the association between the pre-test knowledge score among the community people of regarding farmer's lung with their selected Socio demographic variables.

Hypothesis

1. H_{01} - There is no significant difference between pre-test & post-test knowledge score regarding farmer's lung among the community people
2. H_{02} - There is no significant association between pre-test knowledge score regarding farmer's lung with their selected Socio demographic variables.
3. H_1 - There is a significant difference between pre-test and post-test knowledge score regarding farmer's lung among the community people.
4. H_2 - There is a significant association between pre-test knowledge score regarding farmer's lung with their selected Socio demographic variables.

Methodology

Research Approach: The research approach adopted for the study was Quantitative Evaluative Research Approach.

Research Design: The research design adopted for the study was Quasi Experimental One Group Pre-Test Post-Test Research Design.

Population: The population for the study was community people.

Sample: In this study, the sample was the community people at Ramnagar, Kanpur, UP.

Sample Size: In this study sample size was 200 community people.

Setting of the study: Community people at Ramnagar, Kanpur, UP

Tools and techniques: A structured knowledge questionnaire designed to gauge community members' awareness of Farmer's lung served as the study's instrument. Age, educational attainment, number of children, monthly family income, religion, dietary habits, family type, economic class, farming

experience, and the source of knowledge regarding farmer's lung were among the demographic characteristics that were included in the tool. The instrument was divided into two sections: Section B asked questions pertaining to knowledge, while

Section A dealt with sociodemographic information. A self-administered questionnaire was used for data collection, and both descriptive and inferential statistics were used for analysis.

Result

Table : 01 The association between the pre-test knowledge score among the community people regarding farmer's lung with their selected Socio demographic variables

SL. NO.	DEMOGRAPHIC VARIABLES	X ²	D.F.	INFERENCE
1	Age of the farmer in year			
A	20-29	3.49	6	N.S P>3.49
B	30-39			at 0.05 level
C	40-49			T=12.59
D	>50			
2	Educational status			
A	No formal education	3.9	6	N.S P>3.9
B	Formal education			at 0.05 level
C	Higher secondary education			T=12.59
D	Graduation or above (Professional)			
3	No of children			
A	One	9.91	6	N.S P>9.91
B	Two			at 0.05 level
C	Three			
D	> Three			T=12.59
4	The monthly income of the of the family			
A	<15000/-	30.67	6	S P<30.67
B	15001-25000/-			at 0.05 level
C	25001-35000/-			T=12.59
D	>35001/-			
5	Religion			
A	Hindu	0.207	6	N.S P>0.207
B	Muslim			at 0.05 level
C	Christian			T=12.59
D	Others			
6	Dietary pattern			
A	Vegetarian	0.99	2	N.S

B	Non vegetarian			P>0.99 at 0.05 level T=5.99
7	Types of family			
A	Nuclear	5.79	4	N.S
B	Joint			P>5.79
C	Extended			at 0.05 level T=9.49
8	Economic class of the family			
A	Middle class			S
B	Lower class	16.86	6	P<16.86
C	Lower-middle class			at 0.05 level
D	Upper class			T=12.59
9	Farming Experience			
A	< 3 yr	17.63	6	S
B	3- 5 Yr			P<17.63
C	5-8 Yr			at 0.05 level
D	>8 Yr			T=12.59
10	If farmer having primary information regarding farmer's lunggetting through			
A	Ongoing Programme	53.44	6	S
B	Health Education			P<53.44
C	Mass media			at 0.05 level
D	Family			T=12.59

Comparison of Pre-Test and Post-Test Level of Knowledge Regarding RegardingFarmer's Lung Among the Community People
Table: 02 Comparison of pre-test and post-test level of regarding Farmer's Lung among the community people
n = 200




Figure 01: Pie diagram showing percentage wise distribution of pre-test and post-test level of knowledge regarding farmer's lung among the community people.

Above (Table No. 13, Fig No.10) The pie diagram shows that the mean of pre-test is 12.5 and post-test is 16.5; mean percentage of pre-test is 34% and post-test is 66%; standard deviation of pre-test is 4.7 and post-test is 5.5, mean difference is 4. Thus, it is showing that post-test mean score is higher than pre-test mean score.

Discussion

The discussion chapter shows that after giving the video assisted teaching programme among 200 community members, their knowledge regarding preventive method of farmer's lung was increased than before. In pre-test majority of the sample was having moderate and inadequate knowledge and some were adequate knowledge too. But in post-test, very less participants were having inadequate knowledge and moderate knowledge was increased and adequate knowledge level also increased regarding preventive method of farmer's lung.

Conclusion

The present study was aimed to assess the effectiveness of video assisted teaching programme on knowledge Regarding preventive method of farmer's lung among community members at selected community area of Kanpur Uttar Pradesh. The relevant data was collected and analysed statistically based on the objectives of the study. Following conclusions were drawn. In pre-test knowledge regarding preventive method of farmer's lung, In pre-test out of 200 members (25%) were having inadequate knowledge, (70%) were having moderately adequate knowledge, (5%) were having adequate knowledge in pre-test. While in post-test out of 200 of members of (8%) were having inadequate knowledge, (72%) were having moderately adequate knowledge, (20%) were having adequate knowledge in post-test.

Financial support and sponsorship: Nil

Conflicts of interests: There is no conflict of interest

References

1. Riario Sforza GG, Marinou A. Hypersensitivity pneumonitis: a complex lung disease. *Clin Mol Allergy*. 2017;15:6.
2. Dales RE, Munt PW. Farmer's Lung Disease. *Can Fam Physician*. 1982 Oct;28:1817-20.
3. Deschênes D, Provencher S, Cormier Y. Farmer's lung-induced hypersensitivity pneumonitis complicated by shock. *Respir Care*. 2012 Mar;57(3):464-6.
4. Tao BG, Shen YE, Chen GX, Wu AL, Hong ZR, Yuan JW, Chen BC, Gu XQ. An epidemiological study on farmer's lung among hay grinders in Dafeng County. *Biomed Environ Sci*. 1988 Jun;1(1):13-8.
5. Husman K, Vohlonen I, Terho EO, Mäntyjärvi RA. Precipitins against microbes in mouldy hay in the sera of farmers with farmer's lung or chronic bronchitis and of healthy farmers. *Eur J Respir Dis Suppl*. 1987;152:122-7.
6. Suwalka Deepak, Abbas Aslam, Kamal Mohsin, A Study to Assess the Knowledge of Primipara Mother Regarding Weaning Diet in Selected Rural Area of Kanpur with A View to Develop and Administer Information Booklet https://www.researchgate.net/publication/394849113_GFNPSS-IJMR-6-6-25-05-A-Study-to-Assess-the-Knowledge-of-Primipara-Mother-Regarding-Weaning
7. Lopez M, Salvaggio JE. Epidemiology of hypersensitivity pneumonitis/allergic alveolitis. *Monogr Allergy*. 1987;21:70-86.
8. Cushen B, Sulaiman I, Donoghue N, Langan D, Cahill T, NicDhonncha E, Healy O, Keegan F, Browne M, O'Regan A. High prevalence of obstructive lung disease in non-smoking farmers: The Irish farmers lung health study. *Respir Med*. 2016 Jun;115:13-9.

