ISSN(O):2582-693X
ISSN(P):3139-3314

Review Article

Mental Health Stigma in the Current Scenario: A Review of Social Perceptions, Barriers to Help-Seeking, and Strategies for Stigma Reduction

Manphool Singh[1], Mohammad Hasan[2], Savita Grewal[3], Jyoti[4], Kuruvind Singh Narroka[5], Diwakar Shukla[6]

[1]Vice-Principal Sikar Nursing College, Sikar

[2]Assistant Professor (Mental Health Nursing), Banda Paramedical College and Nursing School Banda(UP)

[3]PhD Scholar, Sharda School of Nursing Sciences and Research, Sharda University, G.D. Uttar Pardesh.

[4]Assistant Professor, V3 College of Nursing, Rudrapur, Uttarakhand

[5]Nursing Tutor (Psychiatric Nursing), College of Nursing, Autonomous State Medical College, Firozabad, U.P.

[6]Principal, Institute of Nursing Research Centre Surajpur Kalua Krishnapur Bharuhamunda Road Surajpur, CG

Corresponding Author:
Dr Manphool Singh, PhD, MSc Nursing in psychiatric,
Vice principal Sikar Nursing College, Sikar

E-mail:
manphool123456@gmail.com

GFNPSS-International Journal of Multidisciplinary Research is a journal of open access. In this journal, we allow all types of articles to be distributed freely and accessible under the terms of the creative common attribution- non-commercial-share. This allows the authors, readers and scholars and general public to read, use and to develop non-commercially work, as long as appropriate credit is given and the newly developed work are licensed with similar terms.

How to cite this article : Singh M, Hasan M, Swami O, Jyoti, Narroka KS, Shukla D. Mental Health Stigma in the Current Scenario: A Review of Social Perceptions, Barriers to Help-Seeking, and Strategies for Stigma Reduction. GFNPSS-IJMR 2026; 7:10: 3747-3751

Submitted: 02 October 2026: **Accepted:** 14 October 2026: **Published:** 31 October 2026

Abstract :

Background: Globally, mental health diseases are a significant public health concern. The stigma attached to mental illness continues to be a major obstacle to early detection, treatment, rehabilitation, and social inclusion despite growing understanding of psychological well-being. People may be discouraged from talking about their mental health issues or getting professional help due to negative attitudes, stereotypes, discrimination, and fear of social rejection.

Keywords: Mental health stigma, mental illness, discrimination, help-seeking behavior, stigma reduction, mental health literacy, and psychological well-being.

INTRODUCTION

A vital part of general health and wellbeing is mental health. It affects people's thoughts, emotions, communication, stress management, interpersonal connections, and social participation. People from a variety of age groups, socioeconomic backgrounds, educational levels, and cultural contexts might be impacted by mental health issues.' However, people with mental health issues may face obstacles that go beyond the signs and symptoms of their illnesses. Stigma is one of the most enduring obstacles.[2]

Negative attitudes, stereotypes, biased views, and discriminatory behaviors aimed against those who suffer from or are thought to suffer from mental health disorders are collectively referred to as mental health stigma.[3] Misunderstandings, fear, false information, cultural attitudes, and the presumption that individuals with mental illness are dangerous, untrustworthy, weak, or incapable of making decisions can all contribute to stigma. The varied experiences and skills of individuals with mental health disorders are not well represented by such presumptions.

Goffman's conceptualization of stigma explains how a person's identity and social relationships can be impacted by socially despised traits. When it comes to mental health, stigma can have an impact on how people view themselves and how society views them.[5] The interrelated processes of labeling, stereotyping, separation, status loss, and discrimination occurring within settings of unequal social power were used by Link and Phelan to further explain stigma.[6]

Despite advancements in digital communication, psychological treatment, public awareness, and mental health education, stigma around mental health is still prevalent today. Discriminatory attitudes have not necessarily vanished despite increased awareness of mental health issues.[7] While acknowledging the significance of mental health in general, people may nonetheless have unfavourable opinions of specific diseases or those who seek psychiatric care.[8]

People may be deterred by stigma from reporting symptoms, talking to family members about emotional challenges, contacting medical specialists, or pursuing therapy.[9] Help-seeking decisions may be further complicated by worries about confidentiality, marital possibilities, career,

educational opportunities, and social reputation. These issues can be particularly significant in societies where mental illness is either poorly understood or closely linked to shame and family honor.[10]

It is important to consider the various cultural, linguistic, religious, economical, and healthcare settings that surround mental health stigma in India. Access to proper care may be delayed by fear of judgment, misconceptions about psychiatric therapy, and worries about social acceptance, even if family ties and community networks may offer crucial emotional and practical support.[11] The accessibility, cost, and availability of mental health care can interact with stigma and affect whether people get help when they need it.[12]

The World Health Organization has emphasized that strategies that uphold human rights and combat prejudice are necessary for mental health promotion, prevention, treatment, and recovery. Reducing the stigma associated with mental illness is therefore not just an educational goal but also a public health, social justice, professional duty, and human dignity issue.[3,4]

This review looks at stigma around mental health in the modern world, focusing on social attitudes, obstacles to getting care, the effects on those who are impacted, and doable tactics to lessen stigma.[8]

SOURCES OF LITERATURE

Databases and academic sites like PubMed, Scopus, Web of Science, PsycINFO, and Google Scholar can be used to find pertinent literature. Reputable publications from the World Health Organization and other reputable public health organizations are also pertinent to this subject.

Key search concepts include:

- Mental health stigma
- Mental illness and discrimination
- Public stigma and self-stigma
- Structural stigma
- Mental health literacy
- Help-seeking behaviour
- Mental health treatment barriers
- Contact-based stigma reduction
- Mental health awareness and education

LITERATURE SELECTION AND INTERPRETATION:

Foundational theoretical papers, pertinent reviews, significant international reports, and research on stigma, help-seeking, discrimination, and stigma-reduction strategies are given priority in the review.

The relationship between personal views, societal expectations, institutional practices, and access to mental health care is taken into consideration while interpreting literature thematically.[1]

ETHICAL CONSIDERATIONS: This article does not entail the recruiting of human subjects or the gathering of new personal health data because it is a literature-based review. The review should appropriately credit its sources and refrain from using language that perpetuates stereotypes or presents individuals with mental health disorders as a monolithic group.[2]

CONCEPT AND TYPES OF MENTAL HEALTH STIGMA: The stigma associated with mental illness is a complex phenomenon that affects social interactions, institutional arrangements, emotional reactions, and beliefs. There may be overlap and reinforcement between several types of stigma.[3]

PUBLIC STIGMA: The term "public stigma" describes the unfavourable opinions, biased attitudes, and discriminatory actions that members of society have against those who suffer from mental health disorders.[4]

Examples include perceiving those receiving psychiatric care as less capable of making judgments, assuming that someone with schizophrenia is always hazardous, or presuming that someone with depression is lazy.[5]

SELF-STIGMA: When people internalize and implement negative societal attitudes about mental illness, it leads to self-stigma. A person may think that having anxiety, sadness, or another mental illness is a sign of failure or weakness. Shame, low self-esteem, hopelessness, and a reluctance to ask for help can all be caused by such ideas.[8]

PERCEIVED STIGMA: The belief or expectation that people with mental health issues will be judged, rejected, or subjected to discrimination by society is known as perceived stigma.[9]

EXPERIENCED STIGMA: Real instances of discrimination, rejection, or unfavourable treatment because of a person's mental health condition are referred to as experienced stigma.[10]

STRUCTURAL STIGMA: Institutional practices, rules, regulations, and social arrangements that disfavour individuals with mental health issues are referred to as structural stigma.[11]

COURTESY STIGMA AND FAMILY STIGMA: When family members, caregivers, or other people connected to a person with mental illness are judged or socially excluded, this is known as courtesy stigma.[12]

MENTAL HEALTH STIGMA IN THE CURRENT SCENARIO

INCREASING AWARENESS AND PERSISTENT MISCONCEPTIONS: Through educational initiatives, public campaigns, online conversations, and increased understanding of psychological well-being in healthcare and educational contexts, mental health awareness has grown.

Even with this advancement, acceptance does not necessarily follow from awareness. People may be aware of typical mental health symptoms, but they may still think that these disorders should be managed on their own without the help of a professional.

INFLUENCE OF CULTURAL AND SOCIAL BELIEFS: People's interpretations of psychological symptoms, odd behaviour, emotional discomfort, and psychiatric therapy are influenced by cultural ideas.

In certain contexts, mental health issues can be solely linked to personal failure, spiritual causes, weak character, or a lack of discipline. Instead of consulting trained medical

https://doi.org/10.46376/ijmr/07.10.2026:3747-3751

professionals, people may initially turn to family members, local authorities, or traditional healers.

SOCIAL MEDIA AND DIGITAL COMMUNICATION: Discussions about anxiety, sadness, stress, trauma, and psychological well-being are now possible thanks to digital media. Online communities can help people find mental health resources and offer peer support.

However, digital communication can also disseminate false information, minimize serious illnesses, encourage self-diagnosis without proper evaluation, or expose people to mockery and cyberbullying.

MENTAL HEALTH STIGMA IN EDUCATIONAL INSTITUTIONS: Academic pressure, exam anxiety, social challenges, bullying, and concern about future career are all possible experiences for students. Despite the growing recognition of the value of psychological well-being by educational institutions, students may be reluctant to seek counselling out of concern that they would be perceived as weak or inadequate.

MENTAL HEALTH STIGMA IN HEALTHCARE SETTINGS: It is expected of healthcare providers to treat patients with compassion and without bias. However, stigma can also arise in healthcare systems when mental health indicators are disregarded, people are held accountable for their illnesses, or psychological issues are given less consideration than physical complaints.

WORKPLACE STIGMA: Discrimination in hiring, a lack of career prospects, exclusion from decision-making, improper disclosure of personal health information, and unfavourable presumptions about dependability and productivity are all examples of workplace stigma.

BARRIERS TO HELP-SEEKING BEHAVIOUR: Recognizing a problem, choosing to ask for help, finding a suitable source of support, and obtaining and continuing care when necessary are all components of help-seeking behaviour.

Every step of this process can be hampered by stigma.

FEAR OF SOCIAL JUDGEMENT: People may be reluctant to talk about emotional problems with friends, family, co-workers, teachers, or medical professionals out of fear of being classified as mentally ill.

LACK OF MENTAL HEALTH KNOWLEDGE: People may be unable to recognize when professional care is necessary due to a lack of information about symptoms, available treatments, and the role of mental health specialists.

CONCERNS ABOUT CONFIDENTIALITY: In environments where people are worried about their social reputation, such as small towns, companies, and educational institutions, confidentiality is especially crucial.

FEAR OF PSYCHIATRIC TREATMENT: Treatment avoidance may be exacerbated by misconceptions regarding hospital-based treatment, counselling, and psychiatric medication.

GENDER AND SOCIAL EXPECTATIONS: Help-seeking behavior may be influenced by societal norms on emotional expression, family responsibilities, masculinity, and femininity.

FAMILY PRESSURE AND COMMUNITY EXPECTATIONS: Because of worries about marriage, work, reputation, or social approval, family members may discourage disclosure.

CONSEQUENCES OF MENTAL HEALTH STIGMA

DELAYED DIAGNOSIS AND TREATMENT: People may hide their symptoms or put off getting a professional evaluation due to stigma. Delaying obtaining assistance might make problems worse and raise the complexity of care needed.

SOCIAL ISOLATION AND REDUCED PARTICIPATION: People who have negative attitudes and a fear of being rejected may avoid relationships, social events, educational chances, and community involvement.

EFFECTS ON SELF-ESTEEM AND PSYCHOLOGICAL WELL-BEING: Recurring discrimination or internalizing unfavourable preconceptions can erode optimism, confidence, and self-worth.

EDUCATIONAL AND OCCUPATIONAL DISADVANTAGES: Bullying, lowered expectations, or exclusion from academic activities are all possible outcomes for students. Workers may experience pressure to hide health issues, prejudice, or limited advancement.

These drawbacks may have an impact on social engagement, career advancement, and financial stability.

IMPACT ON FAMILIES AND CAREGIVERS: Family members may face caregiving obligations, social pressure, emotional suffering, and uncertainty. Families may find it challenging to get information and assistance when stigma prohibits open conversation.

STRATEGIES FOR STIGMA REDUCTION

MENTAL HEALTH EDUCATION AND AWARENESS PROGRAMMES: Understanding of mental health disorders, symptoms, treatment options, and relevant support services can be enhanced through structured educational programs.

Instead than depending solely on generic awareness messages, programs should directly address frequent misconceptions.

CONTACT-BASED INTERVENTIONS: Meaningful engagement with individuals who have personally experienced mental health disorders is a key component of contact-based therapies.

Personal tales, assisted conversations, instructional lectures, and organized interactions between community members and those with lived experience are examples of such interactions.

RESPONSIBLE MEDIA REPRESENTATION: Public perception of mental illness might be influenced by media coverage. Stigma may be strengthened by sensationalized descriptions, false links between mental illness and violence, and disparaging language.

FAMILY AND COMMUNITY PARTICIPATION: Initiatives rooted in the community can promote candid dialog, dispel myths, and assist people in locating suitable resources for assistance.

TRAINING FOR HEALTHCARE PROFESSIONALS: Healthcare workers need to be knowledgeable, skilled communicators, and given the chance to consider their own presumptions.

https://doi.org/10.46376/ijmr/07.10.2026:3747-3751

ANTI-DISCRIMINATION POLICIES AND INSTITUTIONAL REFORM: Alongside educational initiatives, institutional changes are necessary to reduce stigma.

Workplaces, educational institutions, healthcare facilities, and schools should all have explicit policies in place to stop harassment, safeguard private information, offer appropriate assistance, and deal with discriminatory behaviour.

ACCESSIBLE AND AFFORDABLE MENTAL HEALTH SERVICES: When people are unable to get the care they need, awareness campaigns are unlikely to be fully beneficial.

Mental health care must to be reasonably priced, easily accessible, sensitive to cultural differences, and adequately staffed.

PEER SUPPORT AND RECOVERY-ORIENTED CARE: Peer support programs can give people the chance to connect with others who share their worries, learn coping mechanisms, and exchange experiences.

RESPONSIBLE USE OF LANGUAGE: Perceptions of people with mental health issues can be influenced by language.

Healthcare providers, educators, academics, and media outlets should refrain from using disparaging terms and phrases that conflate a diagnosis with an individual's whole identity.

EVALUATION OF STIGMA-REDUCTION INTERVENTIONS: Programs aimed at reducing stigma should be assessed rather than evaluated based only on participation or attendance.**ROLE OF NURSING PROFESSIONALS IN REDUCING MENTAL HEALTH STIGMA:** When it comes to raising awareness of mental health issues and helping people who are experiencing psychological discomfort, nurses play a crucial role. Their duties encompass public health initiatives, educational institutions, hospitals, and community settings.

PATIENT EDUCATION: Using clear language, nurses may clarify common mental health issues, recognize warning signals, give information about available treatments, and dispel myths.

THERAPEUTIC COMMUNICATION: Establishing trust is facilitated by proper confidentiality, empathy, non-judgmental responses, and respectful listening.

EARLY IDENTIFICATION AND REFERRAL: During every day healthcare exchanges, nurses may detect psychological suffering. When necessary, they can assist with evaluation, offer preliminary assistance, and direct them to licensed mental health specialists.

FAMILY COUNSELLING AND SUPPORT: Nurses can assist families in comprehending mental health issues, communicating in a supportive manner, and realizing the significance of receiving the right care.

Family participation should respect the person's autonomy, consent, privacy, and any applicable legal obligations.

ADVOCACY AND PROFESSIONAL ROLE MODELLING: Nurses can take part in institutional mental health efforts, advocate for fair healthcare procedures, and confront discriminatory behaviour.

RESEARCH AND COMMUNITY OUTREACH: Nursing researchers can evaluate educational initiatives, investigate local experiences of stigma, assess mental health literacy, and pinpoint obstacles to service utilization.

IMPLICATIONS FOR EDUCATION, PRACTICE, AND POLICY

IMPLICATIONS FOR NURSING EDUCATION: Nursing courses should incorporate anti-stigma education and mental health literacy. Students should be given the chance to learn about recovery-oriented treatment, reflect on professional attitudes, comprehend human rights, and improve their communication skills.

Classroom instruction may be enhanced by case discussions, supervised clinical experiences, and suitably mediated interactions with those who have lived experience.

CONCLUSION

Psychological well-being, prompt help-seeking, treatment participation, social inclusion, and equitable healthcare are all still significantly hampered by mental health stigma. It shows up as institutional procedures that disadvantage people with mental health disorders, internalized negative views, anticipated judgment, experienced discrimination, and public prejudice.

Providing respectful care, enhancing mental health literacy, enabling early referral, assisting families, and opposing discriminatory behaviors are all crucial tasks for nurses and other medical workers. Policymakers, businesses, educational institutions, and community organizations must all support long-term reform.

FINANCIAL SUPPORT AND SPONSORSHIP: Nil

CONFLICTS OF INTERESTS: There is no conflict of interest

REFERENCES

1. Goffman E. *Stigma: Notes on the Management of Spoiled Identity*. Englewood Cliffs (NJ): Prentice-Hall; 1963.
2. Link BG, Phelan JC. Conceptualizing stigma. Annu Rev Sociol. 2001;27:363-85.
3. World Health Organization. *World Mental Health Report: Transforming Mental Health for All*. Geneva: World Health Organization; 2022.
4. Jorm AF, Korten AE, Jacomb PA, Christensen H, Rodgers B, Pollitt P. "Mental health literacy": a survey of the public's ability to recognise mental disorders and their beliefs about the effectiveness of treatment. Med J Aust. 1997;166(4):182-6.
5. Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rüsch N. Challenging the public stigma of mental illness: a meta-analysis of outcome studies. Psychiatr Serv. 2012;63(10):963-73.
6. Thornicroft G, Rose D, Kassam A, Sartorius N. Stigma: ignorance, prejudice or discrimination? Br J Psychiatry. 2007;190:192-3.
7. Corrigan PW, Watson AC. Understanding the impact of stigma on people with mental illness. World Psychiatry. 2002;1(1):16-20.

https://doi.org/10.46376/ijmr/07.10.2026:3747-3751

8. Clement S, Schauman O, Graham T, Maggioni F, Evans-Lacko S, Bezborodovs N, et al. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. Psychol Med. 2015;45(1):11-27.
9. Henderson C, Evans-Lacko S, Thornicroft G. Mental illness stigma, help seeking, and public health programs. Am J Public Health. 2013;103(5):777-80.
10. Evans-Lacko S, Corker E, Williams P, Henderson C, Thornicroft G. Effect of the Time to Change anti-stigma campaign on trends in mental-illness-related public stigma among the English population in 2003–13: an analysis of survey data. Lancet Psychiatry. 2014;1(2):121-8.
11. Thornicroft G, Mehta N, Clement S, Evans-Lacko S, Doherty M, Rose D, et al. Evidence for effective interventions to reduce mental-health-related stigma and discrimination. Lancet. 2016;387(10023):1123-32.
12. World Health Organization. *Comprehensive Mental Health Action Plan 2013–2030*. Geneva: World Health Organization; 2021.

https://doi.org/10.46376/ijmr/07.10.2026:3747-3751